



Tsinghua Sanya International Mathematics Forum

(清华三亚国际数学论坛)

Future Directions of Photonics and Phononics Workshop

光子和声音的未来方向研讨会

December 11-15, 2017

No.100 Tsinghua Rd., Tianya District, Sanya, Hainan.


(海南省三亚市天涯区清华路100号)

Welcome to TSIMF

The facilities of TSIMF are built on a 23-acre land surrounded by pristine environment at Phoenix Hill of Phoenix Township. The total square footage of all the facilities is over 29,000 square meter that includes state-of-the-art conference facilities (over 10,000 square meter) to hold many international workshops simultaneously, two libraries, a guest house (over 10,000 square meter) and the associated catering facilities, a large swimming pool, gym and sports court and other recreational facilities.

Mathematical Sciences Center (MSC) of Tsinghua University, assisted by TSIMF's International Advisory Committee and Scientific Committee, will take charge of the academic and administrative operation of TSIMF. The mission of TSIMF is to become a base for scientific innovations, and for nurturing of innovative human resource; through the interaction between leading mathematicians and core research groups in pure mathematics, applied mathematics, statistics, theoretical physics, applied physics, theoretical biology and other relating disciplines, TSIMF will provide a platform for exploring new directions, developing new methods, nurturing mathematical talents, and working to raise the level of mathematical research in China.


About Facilities



Registration

Conference booklets, room keys and name badges for all participants will be distributed at the Registry. Please take good care of your name badge. It is also your meal card and entrance ticket for all events.

Guest Room



Conference Center can receive about 378 people having both single and double rooms, and 42 family rooms.

All the rooms are equipped with: free Wi-Fi, TV, air conditioning and other utilities.

Family rooms are also equipped with kitchen and refrigerator.

Library

Opening Hours: 08:00am-22:00pm

TSIMF library is available during the conference and can be accessed by using your room card. There is no need to sign out books but we ask that you kindly return any borrowed books to the book cart in library before your departure.

Restaurant



All the meals are provided in the Chinese Restaurant (Building B1) according to the time schedule.

Breakfast	07:30-08:30
Lunch	12:00-13:30
Dinner	17:30-19:00

Laundry

Opening Hours: 24 hours

The self-service laundry room is located in the Building 1 (B1), next to the shop.


Gym

The gym is located in the Building 1 (B1), opposite to the reception hall. The gym provides various fitness equipment, as well as pool tables, tennis tables and etc.

Playground

Playground is located on the east of the central gate. There you can play basketball, tennis and badminton. Meanwhile, you can borrow table tennis, basketball, tennis balls and badminton at the reception desk.

Swimming Pool



Please note that there are no lifeguards. We will not be responsible for any accidents or injuries. In case of any injury or any other emergency, please call the reception hall at +86-898-38882828.

Outside Shuttle Service:

We have shuttle bus to take participants to the airport for your departure service. Also, we would provide transportation at the Haipo Square (海坡广场) of Howard Johnson for the participants who will stay outside TSIMF. If you have any questions about transportation arrangement, please feel free to contact Ms. Li Ye (叶莉), her cell phone number is (0086)139-7679-8300.

Free Shuttle Bus Service at TSIMF:

We provide free shuttle bus for participants and you are always welcome to take our shuttle bus, all you need to do is wave your hands to stop the bus.

Destinations: Conference Building, Reception Room, Restaurant, Swimming Pool, Hotel etc.



Contact Information of Administration Staffs

Location of Conference Affairs Office: Room 104, Building A

Tel: 0086-898-38263896

Technical Support: Shouxi, He 何守喜

Tel: 0086-186-8980-2225

E-mail: hesx@tsimf.cn

Administrator: Ms. Xianying, Wu 吴显英

Tel: 0086-186-8962-3393

E-mail: wuxy@tsimf.cn

Location of Accommodation Affair Office: Room 200, Building B1

Tel: 0086-898-38882828

Accommodation Manager: Ms. Li Ye 叶莉

Tel: 0086-139-7679-8300

E-mail: yeli@tsimf.cn

Director of TSIMF:

Prof. Xuan Gao 高瑄

Tel: 0086-186-0893-0631

E-mail: gaoxuan@tsinghua.edu.cn

2017 年 12 月 12 日	
Tue. 18:00-20:00 PM	Conference Banquet is on Tuesday night 18:00-20:00 pm
2017 年 12 月 13 日	
Wed. 7:30-8:30 AM	Breakfast
Wed. 8:30-8:50 AM	Group Photo
Wed. 8:50-9:20 AM	Explore Complex Materials with IR and THz near-field Microscopy and Spectroscopy
	<i>Mengkun Liu, Stony Brook University, mengkun@stonybrook.edu</i>
Wed. 9:20-9:50 AM	Mapping three-dimensional near-field responses with scattering-type scanning near-field optical microscopy
	<i>Xiaoji Xu, Lehigh University, xgx214@lehigh.edu</i>
Wed. 9:50-10:10 AM	Coffee Break
Wed. 10:30-11:00 AM	Scattering from singular surfaces, from AFM tip for NSOM to polygonal plates for Terahertz sensor
	<i>S. T. Chui, University of Delaware, chui@bartol.udel.edu</i>
Wed. 11:00-11:30 AM	Low-temperature nano-imaging of polaritons and correlated electrons in quantum materials

	<i>Alexander Swinton McLeod, Columbia University, am4734@columbia.edu</i>
Wed. 11:30-12:00 AM	Optical Study of Topological Crystalline Insulators and Graphene/BN Heterostructures
	<i>Zhiqiang Li, Sichuan University, zliprof@sina.com</i>
Wed. 12:00-13:30 AM	Lunch
Wed. 13:30-17:00 PM	Conference Tour
Wed. 18:00-19:30 PM	Dinner
2017 年 12 月 14 日	
Thu. 8:50-9:20 AM	Synthetic gauge potential in classical waves
	<i>C.T. Chan, Hong Kong University of Science and Technology, phchan@ust.hk</i>
Thu. 9:20-9:50 AM	Observation of optical vortices in momentum space
	<i>Lei Shi, Department of Physics, Fudan University, lshi@fudan.edu.cn</i>
Thu. 9:50-10:10 AM	Coffee Break
Thu. 10:30-11:00 AM	Valley physics in phononic crystals
	<i>Zhengyou Liu, Wuhan University, zyliu@whu.edu.cn</i>
Thu. 11:00-11:30 AM	Manipulating electromagnetic wave propagating non-reciprocally by a chain of magnetic rods

	<i>Rui-Xin Wu, Nanjing University, rxwu@nju.edu.cn</i>
Thu. 11:30-12:00 AM	Anomalous Dipole-like Multipole in Homogenous Dielectric Rods
	<i>JunJie Du, Shanghai Normal University, phyjunjie@gmail.com</i>
2017 年 12 月 15 日	
Fri. 9:00-9:30 AM	Terahertz photonic devices for frequency comb operation and fast detection
	<i>Dr. Wenjian Wan, Shanghai Institute of Microsystem and Information Technology, wjwan@mail.sim.ac.cn</i>
Fri. 9:30-10:00 AM	Quantum behaviors of graphene plasmons
	<i>Changgan Zeng, University of Science and Technology of Chin, cgzeng@ustc.edu.cn</i>
Fri. 10:00-10:30 AM	Coffee Break
Fri. 10:30-11:00 AM	Nonlocal metamaterials
	<i>Yun Lai, Soochow University, laiyun@suda.edu.cn</i>
Fri. 11:00-11:30 AM	Probing optical anisotropy of nanometer-thin van der waals microcrystals by near-field imaging
	<i>Debo Hu, National +A1:B41Center for Nanoscience and Technology, Beijing</i>

Explore Complex Materials with IR and THz near-field Microscopy and Spectroscopy

Prof. Mengkun Liu, Stony Brook University

ABSTRACT: In strongly correlated electron materials (SCEM), the delicate interplay between spin, charge, and lattice degrees of freedom often leads to extremely rich phase diagrams exhibiting intrinsic phase inhomogeneities. The key to studying and disentangling such complexities in SCEM usually lies in the characterization and control of the materials at fundamental energy, time and length scales. I will use this opportunity to report the recent advances in the near-field IR and THz spectroscopy and discuss their applications in the nanoscale electrodynamic in SECM. Specifically, I will discuss the details of modeling scanning near-field infrared microscopy with CST simulation and its application in understanding the mesoscopic insulator to metal phase transitions in VO_2 and Ca_2RuO_4 with over a broad spectral range (350 cm^{-1} to 2500 cm^{-1}). I will also discuss the future development of near-field scanning microscope including the cryogenic capabilities and its coupling to ultrafast pump-probe spectroscopy. These results set the stage for future spectroscopic investigations to access the fundamental properties of complex materials at the nanoscale.

Mengkun Liu (Ph.D. 2012 Boston University) is an assistant professor at the Physics Department of Stony Brook University (since Jan. 2015). His post doc research is at UC San Diego from 2012-2014. His research interests include physics of correlated electron systems, two-dimensional materials, infrared nano-optics and ultrafast time-domain spectroscopy. Prizes include Seaborg Institute Research Fellowships at Los Alamos National Lab (2009, 2010).

Xiaoji Xu, Lehigh University, xgx214@lehigh.edu

Mapping three-dimensional near-field responses with scattering-type scanning near-field optical microscopy"

Scattering-type scanning near-field optical microscopy (s-SNOM) enables mapping of nanoscale field distributions in two dimensions. However, the standard s-SNOM technique lacks direct resolving ability along the vertical direction, therefore unable to provide full three-dimensional near-field responses. Here, we develop a technique that enables s-SNOM to collect near-field responses at explicit tip-sample distance dependences. As a demonstration, spectral responses of boron nitride and silicon carbide are studied by the new s-SNOM method to reveal bound electric near-field of plasmonic structures, tip-induced spectral shift and modified relaxation dynamics of phonon polaritons. The new technique extends the capability of s-SNOM and is generally applicable for a wide range of nanoscale characterizations on plasmonic and phonon polaritonic nanostructures.

Scattering from singular surfaces, from AFM tip for NSOM to polygonal plates for Terahertz sensor

S. T. Chui. Delaware

We study the response of a conical metallic surface to an external electromagnetic (EM) field by representing the fields in basis functions containing integrable singularities at the tip obtained by conformal mapping between the cone and a round disk. We found the field-induced charge distribution to consist of localized terms at the tip apex and the base and an extended bulk term along the body of the cone far way from the tip. We apply our calculation to the scattering-based scanning near-field optical microscope (s-SNOM) and successfully quantify the elastic light scattering from a vibrating metallic tip over a uniform sample. In recent s-SNOM experiments at the optical range ($< 1\mu\text{m}$) the fundamental is much larger than the higher harmonics whereas at THz range ($> 1\text{mm}$) the fundamental becomes comparable to the higher harmonics. We find that the tip charge dominates the contribution to the higher harmonics and becomes much bigger for the THz experiments, thus providing an explanation. Our calculation is orders of magnitude faster than current approaches. We demonstrate this by extracting a two dimensional dielectric constant map from the s-SNOM image of a finite metallic disk which suggests the possibility of mapping the charge density distribution of a finite metallic surface induced by an EM field.

Optical Study of Topological Crystalline Insulators and Graphene/BN Heterostructures

Zhiqiang Li

Sichuan University, zliprof@sina.com

Abstract: The surface states of topological crystalline insulators (TCI) arise from the topology of the bulk bands and are protected by crystalline symmetries. So far, the surface states remain mostly elusive in optical and transport experiments. We investigated the infrared (IR) reflectance spectrum of a TCI $\text{Pb}_{1-x}\text{Sn}_x\text{Se}$ in zero and high magnetic fields. Analysis of the interband Landau level transitions of the TCI phase shows that their bulk bands can be well described by massive Dirac fermions. Moreover, our optical data have revealed several signatures of the surface states in the TCI phase. We will discuss and compare the mobility and DC conductivity of the surface and bulk states. These findings provide new insights in the unique surface states of TCIs.

In the second half of the talk, we report our study of photocurrent in graphene/boron-nitride Moiré superlattice structures. A record-high zero-bias photoresponsivity of 0.3 A/W (equivalently, an external quantum efficiency exceeding 50%) is achieved using graphene's photo-Nernst effect, which demonstrates a collection of at least five carriers per absorbed photon. We reveal that this effect arises from the enhanced Nernst coefficient through Lifshitz transition at low-energy Van Hove singularities, which is an emergent phenomenon due to the formation of Moiré minibands. Our observation points to a new means for extremely efficient and flexible optoelectronics based on van der Waals heterostructures.

Thursday

Synthetic gauge potential in classical waves

C.T. Chan, Hong Kong University of Science and Technology

We show that a synthetic gauge potential in momentum space can be introduced by twisting the micro-structure of a classical wave system in real space. One example is the geometric phase mediated topological transport of sound vortices. It is well-known that geometric phase in light, as a vector wave, manifests through polarization (or spin) that is an intrinsic degree of freedom of vector waves. Such a degree of freedom is absent in scalar waves, such as airborne sound. We demonstrate that the propagation of airborne sound can also induce geometric phase if we create a sound vortex and the geometric phase arises from the propagation of the topological defect state associated with the sound vortex through a helical waveguide in which the 3D twisting bestows a nontrivial winding of the wave vector in momentum space. The effect can be interpreted as a twisting induced synthetic gauge potential that couples orbital angular momentum to the linear momentum. Another example is the control of the equifrequency surfaces at the long wavelength limit. All natural materials have their index ellipsoids centered at $k=0$.

We will see that some metallic scaffolds with helical structures can possess multiple index ellipsoids centered at arbitrary nonzero k -points. The shifting of the equifrequency surfaces at low frequencies can again be interpreted as the consequence of a twisting induced gauge potential. Other examples include the realization of Weyl crystals. Weyl points, as monopoles of Berry curvature in momentum space, have captured much attention recently in various branches of physics. However, many Weyl photonic structures designed previously are probably too complicated for nanofabrication. We found that Weyl points can be found in metallic chiral structures. In the microwave regime, such structures can be fabricated using planar fabrication technology. In IR and optical frequency regimes, Weyl points can be found in chiral woodpile photonic crystals, which can be fabricated using current nanotechnology. When the constituent materials change from the air-in-metal to metal-in-air configuration, the sign of topological charge will change, leading to a topological phase transition and the bands change from topologically trivial to nontrivial. Gapless surface states exist in the latter cases and its robust transport property has been demonstrated.

Observation of optical vortices in momentum space

Yiwen Zhang^{1,4}, Ang Chen^{1,4}, Wenzhe Liu^{1,4}, Chia Wei Hsu³,

Fang Guan^{1,4}, Xiaohan Liu^{1,4}, Lei Shi^{1,4}, Ling Lu² and Jian Zi^{1,4}

1 Department of Physics, Key Laboratory of Micro- and Nano-Photonic Structures (Ministry of Education), and State Key Laboratory of Surface Physics, Fudan University, Shanghai 200433, China

2 Institute of Physics, Chinese Academy of Sciences/Beijing National Laboratory for Condensed Matter Physics, Beijing 100190, China

3 Department of Applied Physics, Yale University, New Haven, Connecticut 06520, USA

4 Collaborative Innovation Center of Advanced Microstructures, Fudan University, Shanghai 200433, China

Email: lshi@fudan.edu.cn

Vortex, the winding of a vector field in two dimensions, has its core the field singularity and its topological charge defined by the quantized winding angle of the vector field [1, 2]. Vortices are one of the most fundamental topological excitations in nature, widely known in hair whorls as the winding of hair strings, in fluid dynamics as the winding of velocities, in angular-momentum beams as the winding of phase angle, and in superconductors and superfluids as the winding of order parameters [2]. Nevertheless, vortices have hardly been observed other than those in the real space. Although band degeneracies, such as Dirac cones, can be viewed as momentum-space vortices in their mathematical structures, there lacks a well-defined physical observable whose winding number is an arbitrary signed integer [3]. Here, we experimentally observed momentum-space vortices as the winding of far-field polarization vectors in the Brillouin zone of periodic plasmonic structures. Using a home-made polarization-resolved momentum-space imaging spectroscopy, we completely map out the dispersion, lifetime and polarization of all radiative states at the visible wavelengths. The momentum space vortices were experimentally identified by their winding patterns in the polarization-resolved iso-frequency contours and their diverging radiative quality factors. Such polarization vortices can exist robustly on any periodic systems of vectorial fields, while they are not captured by the existing topological band theory developed for scalar fields. This work opens up a promising avenue for exploring topological photonics in the momentum space [4], studying bound states in continuum (BICs) [5], as well as for rendering and steering vector beams and designing high-Q plasmonic resonances.

References:

- [1] N. David Mermin, *Rev. Mod. Phys.* 51, 591 (1979)
- [2] J. M. Kosterlitz et al., *J. Phys. C* 5, L124–L126 (1972)
- [3] D. Hsieh et al., *Science* 323, 919–922 (2009).
- [4] L. Lu et al., *Nat. Photon.* 8, 821–829 (2014).
- [5] C. W. Hsu et al., *Nat. Rev. Mater.* 1, 16048 (2016)

Valley physics in phononic crystals

Zhengyou Liu

School of Physics and Technology, Wuhan University

Abstract:

Phononic crystals (PCs) are artificial periodical structures for acoustic waves with frequency bands in momentum space. Dirac conic dispersions can be found at the Brillouin zone of 2D PCs with hexagonal lattice. By breaking the symmetry of the PCs, the degeneracies at two Dirac points can be lifted and bandgaps open, resulting in two inequivalent valleys with states of opposite vortices. The valley vortex states which carry angular momentum can exert torque on particles through interacting with them. Breaking symmetry oppositely leads to two PCs with band inversion, and the one-way edge states emerge on the interfaces of the two PCs, as demonstrated in simulations and in experiments.

Manipulating electromagnetic wave propagating non-reciprocally by a chain of magnetic rods

Rui-Xin Wu

School of Electronic Science and Engineering, Nanjing University, Nanjing 210093, China

Abstract

We theoretically and experimentally demonstrated that non-reciprocal wave propagation could be manipulated by magnetic rod chains under bias DC magnetic fields. We show along the direction of the chain the wave propagation is non-reciprocal, however, the underlying mechanism may be different, One is related to the chiral edge state and the other related to the non-reciprocal bulk state of the rod array. The direction of the wave propagation can be tuned by the DC magnetic field or the symmetry to the chain. The unidirectional transmission makes the chain as one-way waveguide with minimized transvers dimension and flexible geometry. On the other hand, we show out of the chain plane the rod array also exhibits non-reciprocal wave propagation characteristics, showing almost a total reflection when the incident wave obliquely impinged on the rod chain, but nearly a total transmission when the wave reversed its propagation direction. The non-reciprocal wave propagation was directly observed by using the field mapping technique. We show the non-reciprocal wave propagation is due to the non-reciprocal diffraction of the rod chain for the orders 0 and ± 1 . The unique non-reciprocal wave properties, along the magnetic rod chain and out of the rod strip plane, provide new ways to control the flow of EM waves.

JunJie Du,

Shanghai Normal University,

Title: Anomalous Dipole-like Multipole in Homogenous Dielectric Rods

Abstract :

Localized optical scattering behavior might determine the properties of the artificially structured optical materials. Typically, in metamaterials locally resonant units are the basis of metaatoms or metamolecules[1], and in metasurfaces the antenna dispersion and the Pancharatnam-

Berry phase of scatterers are the necessary condition to introduce abrupt phase changes[2]. In this report, we show that circular silicon rods can support an anomalous dipolelike multipole. Similar to an electronic dipole, the radiation fields from the rod are located in the forward and backward directions, but the phase is different from that of an electric dipole. The formation of the dipolelike multipole is also unique, resulting from the interplay between an electric quadrupole and a magnetic dipole excited simultaneously in a dielectric rod. Recently it has been shown that an EM counterpart of anapoles can be similarly obtained by the interplay between a toroidal dipole and an electric dipole. We will show that such silicon rods with the anomalous localized property are able to produce some unexpected results. The compatibility of the designed optical units with complementary metal-oxide semiconductor technologies might make it easily find useful applications in the fields of photonics, optical manipulation.

Friday:

Powerful spintronic THz emitters based on the ultrafast spin-to-charge conversion". I will try to give the abstract asap.

Jingbo Qi, University of Electronic Science and Technology of China

Terahertz photonic devices for frequency comb operation and fast detection

Hua Li

Key Laboratory of Terahertz Solid-State Technology, Shanghai Institute of Microsystem and Information Technology, Chinese Academy of Sciences, 865 Changning Road, Shanghai 200050, China.

** Email: hua.li@mail.sim.ac.cn*

Abstract

The terahertz wave with the frequency defined between 0.1 and 10 THz is of great interest due to its unique properties such as transparency for papers and plastic materials, abundant absorption "fingerprints" of various chemicals, and potential large communication bandwidth. The development of the terahertz technology is strongly dependent on the advances of terahertz sources and detectors. In our laboratory, we have been working on the semiconductor photonic devices employing the intersubband transitions, quantum cascade laser (QCL) and quantum well photodetector (QWP), for terahertz generation and detection, respectively. Here in this work, we report the homogeneous spectral spanning of terahertz QCLs for comb operation and fast terahertz QWP for detection of gigahertz modulated terahertz radiation.

Due to the long cavity geometry that results in a smaller inter-mode beat note frequency and then stronger mode coupling, we can efficiently modulate the long cavity laser and broaden the terahertz emission spectra utilizing a radio frequency modulation technique. The modulated spectra can homogeneously span 330 GHz frequency range, which is approximately 8% of the central frequency. Using the long cavity laser comb with homogeneous spectral spanning, GaAs etalon transmission and ammonia gas (NH_3) absorption measurements are successfully demonstrated.

Concerning the terahertz detection, by carefully designing the device structure and the microwave transmission line, we first demonstrate the fast detection of terahertz light using a terahertz QWP with a device area of $400 \times 400 \mu\text{m}^2$. The electrical rectification measurements reveal that the QWP device has a modulation response bandwidth upto 4.3 GHz. As a proof for the fast detection, a 6.2 GHz modulated terahertz light emitted from the 6-mm long terahertz QCL is successfully detected in terms of inter-mode optical beat note spectrum using the fast terahertz QWP.

References

- [1] G. Scalari, N. Hoyler, M. Giovannini, and J. Faist, "Terahertz bound-to-continuum quantum-cascade lasers based on optical-phonon scattering extraction", *Appl. Phys. Lett.* **86** (18), 181101 (2005).
- [2] M. Wienold, L. Schrottke, M. Giehler, R. Hey, W. Anders, and H. T. Grahn, "Low-threshold terahertz quantum-cascade lasers based on GaAs/Al_{0.25}Ga_{0.75}As heterostructures", *Appl. Phys. Lett.* **97** (7) (2010).
- [3] S. Barbieri, M. Ravaro, P. Gellie, G. Santarelli, C. Manquest, C. Sirtori, S. P. Khanna, E. H. Linfield, and A. G. Davies, "Coherent sampling of active mode-locked terahertz quantum cascade lasers and frequency synthesis", *Nat. Photonics* **5** (5), 306 (2011).
- [4] P. Gellie, S. Barbieri, J. F. Lampin, P. Filloux, C. Manquest, C. Sirtori, I. Sagnes, S. P. Khanna, E. H. Linfield, A. G. Davies, H. Beere, and D. Ritchie, "Injection-locking of terahertz quantum cascade lasers up to 35GHz using RF amplitude modulation", *Opt. Express* **18** (20), 20799 (2010).
- [5] H. Li, P. Laffaille, D. Gacemi, M. Apfel, C. Sirtori, J. Leonardon, G. Santarelli, M. Rösch, G. Scalari, M. Beck, J. Faist, W. Hänsel, R. Holzwarth, and S. Barbieri, "Dynamics of ultra-broadband terahertz quantum cascade lasers for comb operation", *Opt. Express* **23** (26), 33270 (2015).
- [6] W. J. Wan, H. Li, T. Zhou, and J. C. Cao, "Homogeneous spectral spanning of terahertz semiconductor lasers with radio frequency modulation", *Sci. Rep.* **7**, 44109 (2017).
- [7] H. Li, W. J. Wan, Z. Y. Tan, Z. L. Fu, H. X. Wang, T. Zhou, Z. P. Li, C. Wang, X. G. Guo, and J. C. Cao, "6.2-GHz modulated terahertz light detection using fast terahertz quantum well photodetectors", *Sci. Rep.* **7**, 3452 (2017).
- [8] T. Zhou, H. Li, W. J. Wan, Z. L. Fu, and J. C. Cao, Terahertz imaging using photomixers based on quantum well photodetectors, *AIP Advances*, to be published.

Quantum behaviors of graphene plasmons

Changgan Zeng (曾长淦)

Department of Physics/ICQD, University of Science and Technology of China

Discovering exotic quantum effects of plasmons and their coupling with other quasiparticles in solids may help to understand such collective excitations at the fundamental level, and further offer quantum-mechanical solution to design future quantum devices with optimized functionalities. In this talk, I will present our recent progress on the quantum behaviors of plasmons in graphene. In particular, I will show the control of plasmon excitation and propagation by tuning the quantum transmission at Heaviside step-like potentials. Moreover, I will also demonstrate that when the electron-plasmon coupling is introduced, the quantum coherence of electrons in graphene is substantially enhanced.

Nonlocal metamaterials and metasurfaces

Yun Lai

College of Physics, Optoelectronics and Energy & Collaborative Innovation Center of Suzhou
Nano Science and Technology, Soochow University, Suzhou 215006, China

email: laiyun@suda.edu.cn

Abstract – We discuss possible interesting consequences when the material parameters of metamaterials / metasurfaces exhibit wave-vector-dependence, which is also known as nonlocality or spatial dispersion. One interesting possibility is the realization of ultratransparent materials [1], which could reach the ultimate transparency beyond any natural materials. Another interesting consequence is a new approach of cloaking via nonlocal metasurfaces, which may lead to much simpler and thinner cloaking shells than those based on transformation optics. Our work shows great potential in exploring the nonlocal degree of freedoms in metamaterials and metasurfaces.

REFERENCES

1. J. Luo *et al.*, Ultratransparent Media and Transformation Optics with Shifted Spatial Dispersions, *Physical Review Letters* 117, 223901 (2016).