



TSIMF 清华三亚国际数学论坛

数值相对论中的新旧问题

New and old problems in Numerical Relativity

January 6-10, 2025 Room A-121, TSIMF

组织者 ORGANIZERS

Lars Andersson, Beijing Institute of Mathematical Sciences and Applications

Badri Krishnan, Radboud Universiteit

Jörg Frauendiener, University of Otago

Zhoujian Cao(曹周健), Chinese Academy of Sciences

Contents

Contents	2
About	1
Schedule	
January 6, 2025, Monday	2
January 7, 2025, Tuesday	2
January 8, 2025, Wednesday	
January 9, 2025, Thursday	3
January 10, 2025, Friday	4
Titles and Abstracts	
Gungwon Kang	5
Andrzej Rostworowski	5
Chan Park	5
JiaJia Qian	6
Maciej Maliborski	6
Enping Zhou	6
Sizheng Ma	7
Ralf Hiptmair	7
Badri Krishnan	8
Håkan Andreasson	8
Daniel Pook-Kolb	8
Jörg Frauendiener	9
Yuewen Chen	9
Daniela Cors	9
Yu Tian	
Lap Ming Lin	
Kaibo Hu	
Alex Vañó Viñuales	
Welcome to TSIMF	
About Facilities	
Registration	
Guest Room	

Library	
Restaurant	
Laundry	
Swimming Po	اا ا
Free Shuttle	Bus Service at TSIMF
Playground	
Contact Inform	nation of Administration Staff

About

New and old problems in Numerical Relativity 数值相对论中的新旧问题

Date

January 6-10, 2025

Venue

Room A-121, TSIMF

Organizers

Lars Andersson, Beijing Institute of Mathematical Sciences and Applications Badri Krishnan, Radboud Universiteit Jörg Frauendiener, University of Otago Zhoujian Cao(曹周健), Chinese Academy of Sciences

Abstract

The proposed conference will gather leading experts on numerical approaches to the study of the Einstein equations. Recent developments in observations as well as in numerical and computational techniques and perturbation theory motivate a conference that gathers researchers from different communities. The Einstein equations of general relativity play a central role in modern physics. Due to the highly nonlinear

nature of the equations, numerical methods provide an essential tool to explore strong field dynamics. Gravitational wave forms constructed using numerical and perturbative techniques are an essential component in the success of gravitational wave observatories. The planned space based gravitational wave observatories poses new challenges in the analysis of gravitational wave emissions. The large scale dynamics of cosmological models beyond the Newtonian regime is another major challenge at present.

Description of the aim

Our goal is to foster interactions between the different communities doing research on these topics and to stimulate research for attacking major open problems in the field.

Schedule

Time&Date	Monday (Jan. 06)	Tuesday (Jan. 07)	Wednesday (Jan. 08)	Thursday (Jan. 09)	Friday (Jan. 10)
7:30-8:30	Breakfast				
09:00-10:00	Gungwon Kang	Enping Zhou	Håkan Andreasson	Jörg Frauendiener	Kaibo Hu
10:00-10:30	Coffee Break				
10:30-11:30	Andrzej Rostworowski	Sizheng Ma	Daniel Pook-Kolb	Yuewen Chen	Alex Vañó Viñuales
12:00-13:30	Lunch				
14:00-14:30	Chan Park	Ralf Hiptmair		Daniela Cors	
14:30-15:00	JiaJia Qian			Yu Tian	
15:00-15:30	Coffee Break			Coffee Break	
15:30-16:30	Maciej Maliborski	Badri Krishnan		Lap Ming Lin	
16:30-17:30					
17:30-19:00	Dinner				

Schedule

January 6, 2025, Monday

Time 日期	Name 报告人	Title 报告题目	
7:30-8:30	Breakfast		
09:00-10:00	Gungwon Kang	Close black hole encounters and non-merging ringdowns	
10:00-10:30	Coffee Break		
10:30-11:30	Andrzej Rostworowski	Vacuum axisymmetric gravitational collapse, revisited	
12:00-13:30	Lunch		
14:00-14:30	Chan Park	Perturbation of Electromagnetic Waves and Motions of Charged Particles	
14:30-15:00	JiaJia Qian	A discrete de Rham scheme for the exterior calculus Einstein's equations	
15:00-15:30	Coffee Break		
15:30-16:30	Maciej Maliborski	Periodic solutions for simple Hamiltonian PDEs	
16:30-17:30			
17:30-19:00	Dinner		

January 7, 2025, Tuesday

Time 日期	Name 报告人	Title 报告题目	
7:30-8:30		Breakfast	
09:00-10:00	Enping Zhou	Evolution of bare quark stars in numerical relativity	
10:00-10:30		Coffee Break	
10:30-11:30	Sizheng Ma	Cauchy-characteristic matching	
12:00-13:30		Lunch	
14:00-15:00	Ralf Hiptmair	FEEC IN FLOW	
15:00-15:30	Coffee Break		
15:30-16:30	Badri Krishnan	Dynamics of black hole horizons in binary black hole mergers	
16:30-17:30			
17:30-19:00		Dinner	

January 8, 2025, Wednesday

Time 日期	Name 报告人	Title 报告题目
7:30-8:30		Breakfast
09:00-10:00	Håkan Andreasson	Stationary and time dependent solutions of the axially symmetric Einstein-Vlasov system
10:00-10:30		Coffee Break
10:30-11:30	Daniel Pook-Kolb	Quasi-Local Mass in a Binary Black Hole Merger
12:00-13:30		Lunch
14:00-15:00		
15:00-15:30		
15:30-16:30		
16:30-17:30		
17:30-19:00		Dinner

January 9, 2025, Thursday

Time 日期	Name 报告人	Title 报告题目	
7:30-8:30	Breakfast		
09:00-10:00	Jörg Frauendiener	A global approach to the nonlinear perturbation of a black hole by gravitational waves	
10:00-10:30		Coffee Break	
10:30-11:30	Yuewen Chen	On the convergence of the discontinuous Galerkin scheme for Einstein- scalar equations	
12:00-13:30		Lunch	
14:00-14:30	Daniela Cors	Critical collapse simulations with bamps	
14:30-15:00	Yu Tian	Computation meets gravitation: from anti-de Sitter to de Sitter	
15:00-15:30	Coffee Break		
15:30-16:30	Lap Ming Lin	Apparent horizon finders for 3D numerical spacetimes	
16:30-17:30			
17:30-19:00	Dinner		

January 10, 2025, Friday

Time 日期	Name 报告人	Title 报告题目	
7:30-8:30	Breakfast		
09:00-10:00	Kaibo Hu	Differential complexes and compatible finite elements with applications to relativity	
10:00-10:30		Coffee Break	
10:30-11:30	Alex Vañó Viñuales	Hyperboloidal evolution, or how to smoothly reach (future null) infinity	
12:00-13:30		Lunch	
14:00-15:00			
15:00-15:30			
15:30-16:30			
16:30-17:30			
17:30-19:00		Dinner	

Titles and Abstracts

Close black hole encounters and non-merging ringdowns

Gungwon Kang Chung-Ang University

A perturbed black hole produces characteristic radiations whose frequencies possess some physical information of the black hole such as its mass and angular momentum. We have investigated close hyperbolic encounters of two black holes. In addition to the single burst signal associated with the hyperbolic passage, a non-merging ringdown gravitational radiation has been found for the first time. This new type of gravitational waveforms turns out to be due to the tidal deformations of individual constituent black holes through strong gravitational interactions. We briefly report the main results of this phenomenon, and discuss implications on astrophysical observability and testing horizon dynamics in black hole scattering.

Vacuum axisymmetric gravitational collapse, revisited

Andrzej Rostworowski Jagiellonian University

I will discuss my efforts to reproduce the results of the renown Abrahams&Evans (AE) paper on a critical axisymmetric gravitational collapse [Abrahams&Evans, PRL70 2980 (1993)]. There were some such tries in the past but, as far as I know, no one has really succeeded so far. In particular, I will discuss AE ansatz, its linearized approximation, the initial data choice and the bases of the pseudospectral code that I have constructed. I will also try to briefly summarize the efforts undertaken in the last 30 years to repeat AE results and check their conclusions.

Perturbation of Electromagnetic Waves and Motions of Charged Particles

Chan Park

Institute for Gravitational Wave Astronomy, Henan Academy of Sciences

Electromagnetic waves (EMWs) are currently a crucial tool in gravitational wave (GW) observation. Both interferometric GW detectors and pulsar timing arrays observe GWs by measuring perturbations in EMWs. However, these methods are based on geometric optics, which is valid only when the frequency of GWs is much smaller than that of EMWs. Focusing on the fact that EMWs fundamentally represent EM fields, we directly address perturbations in EMWs by obtaining perturbed solutions of Maxwell's equations. To measure the perturbed EMWs, we consider a charge moving within the EMW. A charge moving in a plane EMW has constants of motion, and we analytically derive the perturbations in these constants, discussing the potential of a new method for GW detection based on these findings.

A discrete de Rham scheme for the exterior calculus Einstein's equations

JiaJia Qian Monash University

Applying new numerical techniques to the Einstein's equations is always a challenge due to the complexity and nonlinearity of these equations. In this talk, we propose a discrete de Rham (DDR) scheme for a 3+1 exterior calculus formulation of Einstein's equations. The DDR method works on general polytopal meshes and has an arbitrary order of convergence, as well as reproducing certain aspects of the continuous de Rham complex. For the Maxwell's equations, these properties are enough to achieve a discrete preservation of constraint by reproducing the continuous calculations. For Einstein's equations however, we can preserve certain constraints on the tetrad, but the preservation of the Hamiltonian constraint remains elusive. We present numerical results for two schemes; one based on the simplest coherent set of equations, and the other related to a formulation that we have proved to be well-posed.

Periodic solutions for simple Hamiltonian PDEs

Maciej Maliborski University of Vienna

To better understand the dynamics of asymptotically anti-de Sitter (AdS) spacetimes and the crucial role of time-periodic solutions, we examine simplified models. In this talk, I will present a study of a toy-model Hamiltonian partial differential equation defined on a bounded domain. I will discuss our latest results, highlighting the intricate structure of time-periodic solutions and their implications for the broader question of stability of AdS. This is joint work with Filip Ficek.

Evolution of bare quark stars in numerical relativity

Enping Zhou Huazhong University of Science and Technolog

Based on the Witten's Conjecture, i.e., 3-flavor quark matter might be the stable state of dense matter, it is believed that at least part of the conventional 'neutron stars' are actually quark stars, if not all of them. Unlike neutron stars, the surface density of quark stars are extremely large (and hence they are called bare quark stars) and comparable to nuclear saturation density. Such finite surface density results in difficulties to model the them numerically. Nevertheless, it is essential to model processes such as the merger of quark stars in order to understand the differences between them and binary neutron star mergers and to distinguish them by observations. In this talk, I will introduce our attempts to model bare quark stars in numerical relativity by modifying the primitive recovery procedure accordingly. I will also introduce the differences between binary quark star merger and binary neutron star merger we found, in the collapse behavior and matter ejections during the post-merger phase.

Cauchy-characteristic matching

Sizheng Ma Perimeter Institute

Cauchy-characteristic matching is a novel numerical-relativity technique that enables the evolution of Einstein's equations on an effectively infinite computational domain. In this talk, I will present our recent successful applications of this method to binary black hole mergers, and demonstrate its transformative potential for modeling tail and memory effects in eccentric binaries. I will also discuss my latest work on extending Cauchy-characteristic extraction (and matching) to theories beyond General Relativity.

FEEC IN FLOW

Ralf Hiptmair ETH Zurich

Speaker: R. HIPTMAIR (Seminar for Applied Mathematics, ETH Zürich, joint work with H. Heumann, C. Pagliantini, and W. Tonnon)

I am going to review mesh-based Eulerian and Lagrangian discretizations of transient and stationary advection-diffusion boundary value problems (BVPs) for differential forms. Advection is passive, assuming the underlying velocity vector field to be given and sufficiently regular.

Though this subject has been exhaustively studied for 0-forms, the development of numerical methods for higher degrees started only 15 years ago, guided by the principle that numerical methods designed for the case of 0-forms can usually be generalized to ℓ -forms with suitable adaptations. As in the case of 0-forms the main challenge is stabilization by taking into account the direction of the flow.

The Eulerian methods start from (localized) standard variational formulations of the advectiondiffusion BVP, restrict them to conforming or broken (in the case of discontinuous Galerkin (DG) methods) trial and test spaces, and add stabilization terms. Such stabilized Galerkin methods come in various flavors: upwind DG, Galerkin least squares, conforming interior penalty, upwind projection, to name some of them. The Lagrangian methods directly discretize the material derivative by tracking the flow. We discuss semi-Lagrangian discretizations on a fixed mesh and will give a glimpse of their application to non-linear transport. References:

[1] H. HEUMANN AND R. HIPTMAIR, Convergence of lowest order semi-Lagrangian schemes, Foundations of Computational Mathematics, 13 (2013), pp. 187–220.

[2] H. HEUMANN AND R. HIPTMAIR, Stabilized Galerkin methods for magnetic advection, ESAIM Math. Model. Numer. Anal., 47 (2013), pp. 1713–1732.

[3] H. HEUMANN, R. HIPTMAIR, K. L I, AND J. XU, Fully discrete semi-Lagrangian methods for advection of differential forms, BIT Numerical Mathematics, 52 (2012), pp. 981–1007.

[4] H. HEUMANN, R. HIPTMAIR, AND C. PAGLIANTINI, Stabilized Galerkin for transient advection of differential forms, Discrete and Continuous Dynamical Systems - Series S, 9 (2016), pp. 185–214.

[5] R. HIPTMAIR AND C. PAGLIANTINI, Splitting-based structure preserving discretizations for magnetohydrodynamics, SMAI J. Comput. Math., 4 (2018), pp. 225–257.

[6] W. TONNON AND R. HIPTMAIR, Semi-Lagrangian finite element exterior calculus for

incompressible flows, Adv. Comput. Math., 50 (2024), pp. Paper No. 11, 29.

TBA

Badri Krishnan Radboud University, Nijmegen, The Netherlands

TBA

Stationary and time dependent solutions of the axially symmetric Einstein-Vlasov system.

Håkan Andreasson University of Gothenburg

In this talk I will present a numerical method used to obtain axially symmetric stationary solutions of the Einstein-Vlasov system. It is based on a mass-preserving algorithm which has been used successfully to construct a wide range of stationary solutions. Recently it was shown that this algorithm converges in a simple case. A few numerical results and open problems will be discussed, in particular those that are related to highly compact solutions. I will also consider the evolution problem and discuss methods such as the particle in cell method and the finite volume method. A few examples related to the weak cosmic censorship conjecture and the Hoop conjecture will be shown.

Quasi-Local Mass in a Binary Black Hole Merger

Daniel Pook-Kolb

Max Planck Institute for Gravitational Physics/Leibniz University Hannover

One of the major open problems in classical general relativity is how one should define the mass of a finite region of space. In this talk, we will investigate a promising definition proposed by Wang and Yau in 2009. A closed 2-surface bounding the region of interest is embedded isometrically into Minkowski space. The mass is then calculated by comparing the extrinsic geometries. The Wang-Yau mass has many desirable properties, but it has previously not been calculated for surfaces in dynamical spacetimes. To remedy this, we will discuss how the Wang-Yau mass can be computed in practice in numerical simulations and also extend the original definition to surfaces important in black hole dynamics: their quasi-local horizons. Finally, we look at how this mass behaves in a merger of black holes.

A global approach to the nonlinear perturbation of a black hole by gravitational waves

Jörg Frauendiener University of Otago

It is well known that gravitational waves interact in a non-linear way. This makes it difficult to describe them rigorously. The cleanest description is based on certain conformal properties of the Einstein equations — first discovered by R. Penrose they were rigorously developed and used by H. Friedrich to prove several important global results for general relativistic space-times. The conformal field equations which implement this conformal framework provide various well-posed initial (boundary) value problems for use in many different situations. The talk will give a computational perspective on one particular application, the non-linear interaction of gravitational waves with an initially static (and spherically symmetric) black hole. We will show how to 'kick' the black hole and possibly how to spin it up.

On the convergence of the discontinuous Galerkin scheme for Einstein-scalar equations

Yuewen Chen Tsinghua University

We prove the stability and convergence of the high order discontinuous Galerkin scheme to spherically symmetric Einstein-scalar equations for a class of large initial data that ensures the formation of a black hole. Having chosen the Bondi coordinate system, we achieve L^2 stability and obtain the optimal error estimates. This is a collaborative work with Professor Chi-Wang Shu and Professor Shing-Tung Yau.

Critical collapse simulations with bamps

Daniela Cors University of Cambridge

Reliable numerical simulations require a thorough control of all non-physical dynamics that can appear, such as constraint violations and coordinate singularities. These become more challenging in the strong-field regime and particularly near the threshold of collapse, where, in the absence of a horizon, arbitrarily large curvatures cannot be excised. At this threshold, a regular and universal structure emerges in spherical symmetry, referred to as critical phenomena. Beyond sphericity, the picture of critical collapse is unclear. Clarification demands simulations that are closer to the threshold. To that end, we have proposed formulation improvements for critical collapse simulations. In this talk I will present a modified generalised-harmonic damping scheme and a self-similarity compatible gauge designed to tune closer to the threshold. I will also show what we have been able to learn about criticality beyond spherical symmetry thanks to these improvements.

Computation meets gravitation: from anti-de Sitter to de Sitter

Yu Tian

University of Chinese Academy of Sciences

Dynamics related to gravity and its applications, including but not limited to gravitational waves, are important in various fields in theoretical physics, which inevitably need numerics. This talk will focus on the formalisms and applications of numerical relativity in asymptotic anti-de Sitter space-times (related to AdS/CFT) and asymptotic de Sitter space-times.

Apparent horizon finders for 3D numerical spacetimes

Lap Ming Lin The Chinese University of Hong Kong

Apparent horizon is an important concept for characterizing the existence and properties of black holes in numerical spacetimes. Apparent horizon finders for generic 3D spacetimes have been developed using different techniques. In this talk, we shall give a brief review of the topic and present our latest apparent horizon finder based on the multigrid method. Our finder has been tested with both analytic and numerical data, including dynamically inspiraling binary black hole spacetimes. Our multigrid-based finder performs comparably to AHFinderDirect, the current fastest open-source finder, in terms of accuracy and starts to outperform it at high angular resolutions in terms of speed. This suggests that the multigrid algorithm provides an alternative option for studying apparent horizons, especially when high resolutions are needed.

Differential complexes and compatible finite elements with applications to relativity

Kaibo Hu The University of Edinburgh

Differential complexes are sequences of spaces connected by differential operators. In this talk, we discuss differential complexes of tensors (hyperforms), generalizing the de Rham sequences and encoding differential structures of tensor-valued problems from continuum mechanics, differential geometry and general relativity etc. Associated with each complex, various PDE problems are formulated, such as the Hodge Laplacian, Hodge Dirac and Hodge wave problems. We discuss possibilities of reformulating the linearized Einstein equations in these forms.

The Finite Element Exterior Calculus (FEEC) provides a framework for discretizing formulations based on complexes. In particular, the de Rham complex enjoys a canonical finite element discretization, unified as finite element differential forms. We review the construction and discuss process in discretizing tensor-valued spaces and complexes.

Hyperboloidal evolution, or how to smoothly reach (future null) infinity

Alex Vañó Viñuales University of Lisbon

Gravitational wave radiation is only unambiguously defined at future null infinity, the collection of the end-points of future-directed null geodesics. A convenient way to reach null infinity in numerical relativity simulations is via hyperboloidal foliations: smooth spacelike slices suitable for solving the Einstein equations as an initial value problem. Spatial compactification on a hyperboloidal slice ensures that future null infinity is included in the numerical integration domain at finite coordinate distance. However, it also requires a regularization of the Einstein equations, and how this is achieved gives rise to different approaches to hyperboloidal evolution. I will give an overview of past and present implementations of the hyperboloidal method to general relativity, putting special emphasis on those being actively developed, as well as report on current progress.



The facilities of TSIMF are built on a 23-acre land surrounded by pristine environment at Phoenix Hill of Phoenix Township. The total square footage of all the facilities is over 29,000 square meter that includes state-of-the-art conference facilities (over 10,000 square meter) to hold many international workshops simultaneously, two reading rooms of library, a guest house (over 10,000 square meter) and the associated catering facilities, a large swimming pool, gym and sports court and other recreational facilities.

Management Center of Tsinghua Sanya International Forum is responsible for the construction, operation, management and service of TSIMF. The mission of TSIMF is to become a base for scientific innovations, and for nurturing of innovative human resource; through the interaction between leading mathematicians and core research groups in pure mathematics, applied mathematics, statistics, theoretical physics, applied physics, theoretical biology and other relating disciplines, TSIMF will provide a platform for exploring new directions, developing new methods, nurturing mathematical talents, and working to raise the level of mathematical research in China.

园区平面图 Park Plan 客房服务中心 Service Center ^{前台电话} Front Desk No. 餐厅 Restaurant 便利店 洗衣房 数学大师园 Master Garden 数学家小道 Math Trail Laund N 泳池 清华路 Tsinghua Road 健身房 ^{Gym} 园区大门 会议大楼 Park Entrance 运动场 Playground Confe 巴士乘车处 停车场 Parking Lot **Bus Stand**

About Facilities

Registration

Conference booklets, room keys and name badges for all participants will be distributed at the front desk. Please take good care of your name badge. It is also your meal card and entrance ticket for all events.



Guest Room

All the rooms are equipped with: free Wi-Fi (Password:tsimf123), TV, air conditioning and other utilities.

Family rooms are also equipped with kitchen and refrigerator.





Library

Opening Hours: 09:00am-22:00pm

TSIMF library is available during the conference and can be accessed by using your room card. There is no need to sign out books but we ask that you kindly return any borrowed books to the book cart in library before your departure.



In order to give readers a better understanding of the contributions made by the Fields Medalists, the library of Tsinghua Sanya International Mathematics Forum (TSIMF) instituted the Special Collection of Fields Medalists as permanent collection of the library to serve the mathematical researchers and readers.

So far, there are 271 books from 49 authors in the Special Collection of Fields Medalists of TSIMF library. They are on display in room A220. The participants are welcome to visit.



Breakfast 07:30-08:45 Lunch 12:00-13:30 Dinner 17:30-19:00

Restaurant

All the meals are provided in the restaurant (Building B1) according to the time schedule.





Laundry

Opening Hours: 24 hours

The self-service laundry room is located in the Building(B1).



Gym

Opening Hours: 24 hours

The gym is located in the Building 1 (B1), opposite to the reception hall. The gym provides various fitness equipment, as well as pool tables, tennis tables etc.



Playground

Playground is located on the east of the central gate. There you can play basketball, tennis and badminton. Meanwhile, you can borrow table tennis, basketball, tennis balls and badminton at the reception desk.

Swimming Pool

Please enter the pool during the open hours, swimming attire and swim caps are required, if you feel unwell while swimming, please stop swimming immediately and get out of the pool. The depth of the pool is 1.2M-1.8M. Opening Hours: 13:00-14:00 18:00-21:00



Free Shuttle Bus Service at TSIMF

We provide free shuttle bus for participants and you are always welcome to take our shuttle bus, all you need to do is wave your hands to stop the bus.

Destinations: Conference Building, Reception Room, Restaurant, Swimming Pool, Hotel etc.



Contact Information of Administration Staff

Location of Conference Affairs Office: Room 104, Building A

Tel: 0086-898-38263896 Conference Affairs: Shouxi He 何守喜 Tel:0086-186-8980-2225 Email: heshouxi@tsinghua.edu.cn

Location of Accommodation Affairs Office: Room 200, Building B1

Tel: 0086-898-38882828 Accommodation Manager: Ms. Li YE 叶莉 Tel: 0086-139-7679-8300 Email: yel@tsinghua.edu.cn

IT

Yuanhang Zhou 周远航 Tel: 0086-133-6898-0169 Email: 13368980169@163.com

*Reception duty hours: 7:00-23:00, chamber service please call: 0086-38882828 (exterior line)
80000 (internal line)
*Room maintainer night duty hours: 23:00-7:00, if you need maintenance services, please call: 0086-38263909 (exterior line) 30162 (internal line)

Director Assistant of TSIMF

Kai CUI 崔凯 Tel/Wechat: 0086- 136-1120-7077 Email :cuik@tsinghua.edu.cn

Director of TSIMF

Prof.Xuan GAO 高瑄 Tel: 0086-186-0893-0631 Email: gaoxuan@tsinghua.edu.cn 清华大学三亚国际论坛管理中心 | 三亚清华数学论坛管理中心 Tsinghua Sanya International Mathematics Forum (TSIMF)

- 0086-898-38882828 0086-898-38883896 0086-898-38883895
- tsimf@tsinghua.edu.cn

572000

https://www.tsimf.cn

海南省三亚市天涯区清华路 100 号,清华三亚国际数学论坛 No.100, Tsinghua Road, Tianya District, Sanya, Hainan, P. R. China.

